

# Fonctions préharmoniques et applications conformes

KILIAN RASCHEL



Journée en hommage à Jacques Neveu  
Institut Henri Poincaré  
23 mai 2017

Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant


## Introductory example & definition

### Markov chains: example



## Introductory example & definition

### Markov chains: example




$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$

$$\text{Solution: } f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1$$

## Introductory example & definition

### Markov chains: example



$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$


$$\text{Solution: } f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1$$

### Markov chains: general theorem

The hitting probabilities are characterized as being the *minimal non-negative solutions* to a system of *linear recurrences*.

## Introductory example & definition

### Markov chains: example



$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$

$$\text{Solution: } f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1$$

### Markov chains: general theorem

The hitting probabilities are characterized as being the *minimal non-negative solutions* to a system of *linear recurrences*.


**Definition:**  $f$  harmonic if  $L[f](x) = 0$  for all  $x$  in a region  $\subset \mathbb{Z}^d$

$$L[f](x) = \sum_{y \in N_y} p(y) \{f(x+y) - f(x)\},$$

with *sets of neighbors*  $N_y \subset \mathbb{Z}^d$  and *weights*  $p = \{p(y)\}_{y \in \mathbb{Z}^d}$

## Introductory example & definition

### Markov chains: example



$$f_i = \mathbb{P}_i[\text{hit } 4] \text{ satisfies } \begin{cases} f_1 = 0 \\ f_4 = 1 \\ f_2 = \frac{1}{2}f_1 + \frac{1}{2}f_3 \\ f_3 = \frac{1}{2}f_2 + \frac{1}{2}f_4 \end{cases}$$

$$\text{Solution: } f_1 = 0, f_2 = \frac{1}{3}, f_3 = \frac{2}{3}, f_4 = 1$$

### Markov chains: general theorem

The hitting probabilities are characterized as being the *minimal non-negative solutions* to a system of *linear recurrences*.

**Definition:**  $f$  harmonic if  $L[f](x) = 0$  for all  $x$  in a region  $\subset \mathbb{Z}^d$

$$L[f](x) = \sum_{y \in N_y} p(y) \{f(x+y) - f(x)\},$$

with *sets of neighbors*  $N_y \subset \mathbb{Z}^d$  and *weights*  $p = \{p(y)\}_{y \in \mathbb{Z}^d}$

▷ Multivariate linear recurrences with constant coefficients

## {History of/Questions on} preharmonic functions (1/2)

### Classical (continuous) harmonic functions in $\mathbb{R}^d$

$$\Delta[f](x) = \sum_{i=1}^d \frac{\partial^2 f(x)}{\partial x_i^2} = 0$$

- ▷ Possibility of adding *weights*  $\leadsto$  *elliptic operators*
- ▷ *Harmonic functions satisfy various properties*: maximum principle/mean value property/Harnack inequalities/Liouville's theorem/relations with analytic functions/etc.
- ▷ *Examples of application*: Heat equation/Dirichlet problem/Poisson's equation/more general PDEs/etc.







## {History of/Questions on} preharmonic functions (1/2)

### Classical (continuous) harmonic functions in $\mathbb{R}^d$

$$\Delta[f](x) = \sum_{i=1}^d \frac{\partial^2 f(x)}{\partial x_i^2} = 0$$




- ▷ Possibility of adding *weights*  $\leadsto$  *elliptic operators*
- ▷ *Harmonic functions satisfy various properties*: maximum principle/mean value property/Harnack inequalities/Liouville's theorem/relations with analytic functions/etc.
- ▷ *Examples of application*: Heat equation/Dirichlet problem/Poisson's equation/more general PDEs/etc.

### Do preharmonic functions satisfy similar properties?

- ▷ Dirichlet problem  Phillips & Wiener '23; Bouligand '25
- ▷ Harnack inequalities  Lawler & Polaski '92; Varopoulos '99
- ▷ Maximum principle, Liouville's theorem & related topics  Heilbronn '48
- ▷ Cauchy-Riemann equations  Duffin '55; Kiselman '05-'08




## {History of/Questions on} preharmonic functions (2/2)

### Further properties


- ▷ Rate of growth  Murdoch '63-'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63-'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

## {History of/Questions on} preharmonic functions (2/2)

### Further properties




- ▷ Rate of growth  Murdoch '63-'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63-'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

### Preharmonic & harmonic functions

- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

## {History of/Questions on} preharmonic functions (2/2)



### Further properties

- ▷ Rate of growth  Murdoch '63–'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63–'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

### Preharmonic & harmonic functions




- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

### Probability theory models

- ▷ Ising models  Mercat '01; Smirnov '10; Chelkak '11; Beffara '12
- ▷ Conformal invariance of lattice models  Duminil-Copin & Smirnov '12

## {History of/Questions on} preharmonic functions (2/2)



### Further properties

- ▷ Rate of growth  Murdoch '63–'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63–'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

### Preharmonic & harmonic functions

- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

### Probability theory models




- ▷ Ising models  Mercat '01; Smirnov '10; Chelkak '11; Beffara '12
- ▷ Conformal invariance of lattice models  Duminil-Copin & Smirnov '12

### Special discrete functions

- ▷ Conformal mappings  Ferrand '44; Isaacs '52
- ▷ Discrete harmonic polynomials & discrete exponential functions  Terracini '45–'46; Heilbronn '48; Isaacs '52; Duffin '55; Duffin & Peterson '68

## {History of/Questions on} preharmonic functions (2/2)



### Further properties

- ▷ Rate of growth  Murdoch '63–'65; Ignatiuk-Robert '10
- ▷ Picard's theorem (sign of harmonic functions) & factorization  Murdoch '63–'65
- ▷ Absolute monotonicity  Lippner & Mangoubi '15

### Preharmonic & harmonic functions

- ▷ Relations between discrete & continuous harmonic functions  Lusternik '26; Ferrand '44; Kesten '91; Varopoulos '09

### Probability theory models

- ▷ Ising models  Mercat '01; Smirnov '10; Chelkak '11; Beffara '12
- ▷ Conformal invariance of lattice models  Duminil-Copin & Smirnov '12

### Special discrete functions


- ▷ Conformal mappings  Ferrand '44; Isaacs '52
- ▷ Discrete harmonic polynomials & discrete exponential functions  Terracini '45–'46; Heilbronn '48; Isaacs '52; Duffin '55; Duffin & Peterson '68

### Potential theory

- ▷ Martin boundary  Woess '92; Kurkova & Malyshev '98; Ignatiuk-Robert & Loree '10; Mustapha '15

## Warning: lattice walk enum. vs. preharmonic functions


Multivariate recurrence relations in both cases



- ▷  $q(n; i, j) = \#_{\mathbb{N}^2} \{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷  $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$   
(Caloric functions)

## Warning: lattice walk enum. vs. preharmonic functions

Multivariate recurrence relations in both cases



- ▷  $q(n; i, j) = \#\_{\mathbb{N}^2}\{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷  $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$   
(Caloric functions)
- ▷  $f(i, j) = \frac{1}{4}\{f(i-1, j) + f(i+1, j) + f(i, j-1) + f(i, j+1)\}$   
(Preharmonic functions)


## Main differences & difficulties

- ▷ A unique solution vs. an unknown ( $\leq \infty$ ) number of solutions
- ▷ Consequence: *guess and prove* techniques do not work



## Warning: lattice walk enum. vs. preharmonic functions

### Multivariate recurrence relations in both cases



- ▷  $q(n; i, j) = \#_{\mathbb{N}^2} \{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷  $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$   
(Caloric functions)
- ▷  $f(i, j) = \frac{1}{4} \{f(i-1, j) + f(i+1, j) + f(i, j-1) + f(i, j+1)\}$   
(Preharmonic functions)

### Main differences & difficulties


- ▷ A unique solution vs. an unknown ( $\leq \infty$ ) number of solutions
- ▷ Consequence: *guess and prove* techniques do not work
- ▷ Generating functions of preharmonic functions satisfy *kernel functional equations*
- ▷ Preharmonic functions  $\approx$  homogenized enumeration problem:

$$K(x, y)Q(x, y) = K(x, 0)Q(x, 0) + K(0, y)Q(0, y) - K(0, 0)Q(0, 0) - xy$$

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

## Warning: lattice walk enum. vs. preharmonic functions

### Multivariate recurrence relations in both cases



- ▷  $q(n; i, j) = \#_{\mathbb{N}^2} \{(0, 0) \xrightarrow{n} (i, j)\}$
- ▷  $q(n+1; i, j) = q(n; i-1, j) + q(n; i+1, j) + q(n; i, j-1) + q(n; i, j+1)$   
(Caloric functions)
- ▷  $f(i, j) = \frac{1}{4} \{f(i-1, j) + f(i+1, j) + f(i, j-1) + f(i, j+1)\}$   
(Preharmonic functions)

### Main differences & difficulties

- ▷ A unique solution vs. an unknown ( $\leq \infty$ ) number of solutions
- ▷ Consequence: *guess and prove* techniques do not work
- ▷ Generating functions of preharmonic functions satisfy *kernel functional equations*
- ▷ Preharmonic functions  $\approx$  homogenized enumeration problem:  
$$K(x, y)Q(x, y) = K(x, 0)Q(x, 0) + K(0, y)Q(0, y) - K(0, 0)Q(0, 0) - xy$$
$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$
- ▷ Preharmonic functions  $\rightsquigarrow$  counting numbers asymptotics

Introduction & motivations

**Applications in probability theory**


Applications in combinatorics

Discrete harmonic functions in the quadrant

## Doob transform

(1/2)


**Example:** construct a 1D process conditioned to stay in  $\mathbb{N}$



## Doob transform


(1/2)

**Example:** construct a 1D process conditioned to stay in  $\mathbb{N}$




- ▷ Function  $f(i) = i$  is positive harmonic and  $f(0) = 0$
- ▷ Weights  $p(i, i \pm 1) = \frac{1}{2}$  become  $p^f(i, i \pm 1) = \frac{1}{2} \frac{f(i \pm 1)}{f(i)} = \frac{1}{2} \frac{i \pm 1}{i}$

**Example:** construct a 1D process conditioned to stay in  $\mathbb{N}$




- ▷ Function  $f(i) = i$  is positive harmonic and  $f(0) = 0$
- ▷ Weights  $p(i, i \pm 1) = \frac{1}{2}$  become  $p^f(i, i \pm 1) = \frac{1}{2} \frac{f(i \pm 1)}{f(i)} = \frac{1}{2} \frac{i \pm 1}{i}$
- ▷ New weights sum to 1:  $f(i-1) + f(i+1) = 2f(i)$




- ▷ Discrete Bessel process

 Biane '90; Mishchenko '05

**Example:** construct a 1D process conditioned to stay in  $\mathbb{N}$



- ▷ Function  $f(i) = i$  is positive harmonic and  $f(0) = 0$
- ▷ Weights  $p(i, i \pm 1) = \frac{1}{2}$  become  $p^f(i, i \pm 1) = \frac{1}{2} \frac{f(i \pm 1)}{f(i)} = \frac{1}{2} \frac{i \pm 1}{i}$
- ▷ New weights sum to 1:  $f(i-1) + f(i+1) = 2f(i)$



- ▷ Discrete Bessel process 📖 Biane '90; Mishchenko '05

## Construction can be generalized

- ▷ *Random processes* conditioned never to leave *cones* of  $\mathbb{Z}^d$
- ▷ Quantum random walks, eigenvalues of random matrices, non-colliding random walks, etc.

📖 Dyson '62; Biane '90-'92; Eichelsbacher & König '08

**A second way for the conditioning** (with the first exit time)

▷ RW  $\{S(n)\}_{n \geq 0}$ ; *first exit time*  $\tau = \inf\{n \geq 0 : S(n) = 0\}$



## A second way for the conditioning (with the first exit time)

- ▷ RW  $\{S(n)\}_{n \geq 0}$ ; *first exit time*  $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event  $\{\tau = \infty\}$ , the RW stays in  $\{1, 2, 3, \dots\}$

**A second way for the conditioning** (with the first exit time)

- ▷ RW  $\{S(n)\}_{n \geq 0}$ ; *first exit time*  $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event  $\{\tau = \infty\}$ , the RW stays in  $\{1, 2, 3, \dots\}$
- ▷ Replace  $\mathbb{P}[\cdot]$  by  $\mathbb{P}[\cdot | \{\tau = \infty\}]$  to obtain a *conditioned RW*


**A second way for the conditioning** (with the first exit time)

- ▷ RW  $\{S(n)\}_{n \geq 0}$ ; *first exit time*  $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event  $\{\tau = \infty\}$ , the RW stays in  $\{1, 2, 3, \dots\}$
- ▷ Replace  $\mathbb{P}[\cdot]$  by  $\mathbb{P}[\cdot | \{\tau = \infty\}]$  to obtain a *conditioned RW*
- ▷ Important question: do we have  $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$ ?
  - ✎ Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time  $\tau$

### A second way for the conditioning (with the first exit time)

- ▷ RW  $\{S(n)\}_{n \geq 0}$ ; *first exit time*  $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event  $\{\tau = \infty\}$ , the RW stays in  $\{1, 2, 3, \dots\}$
- ▷ Replace  $\mathbb{P}[\cdot]$  by  $\mathbb{P}[\cdot | \{\tau = \infty\}]$  to obtain a *conditioned RW*
- ▷ Important question: do we have  $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$ ?
  - 📎 Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time  $\tau$

### Example (1/3) in the quadrant: the simple walk




- ▷ Uniform weights  $\frac{1}{4}$
- ▷  $f(i, j) = i \cdot j$
- ▷ *Unique preharmonic function* (up to multiplicative factors)
- ▷ Product form 📎 Picardello & Woess '92



### A second way for the conditioning (with the first exit time)

- ▷ RW  $\{S(n)\}_{n \geq 0}$ ; *first exit time*  $\tau = \inf\{n \geq 0 : S(n) = 0\}$
- ▷ On the event  $\{\tau = \infty\}$ , the RW stays in  $\{1, 2, 3, \dots\}$
- ▷ Replace  $\mathbb{P}[\cdot]$  by  $\mathbb{P}[\cdot | \{\tau = \infty\}]$  to obtain a *conditioned RW*
- ▷ Important question: do we have  $\mathbb{P}[\cdot | \{\tau = \infty\}] = \mathbb{P}^f$ ?
  - 📎 Denisov & Wachtel '15 (general cones with zero drift); Courtiel, Melczer, Mishna & R. '16 (Gouyou-Beauchamps model with drift)
- ▷ Difficulty: exit time  $\tau$

### Example (3/3) in the quadrant: the GB walk



- ▷ Uniform weights  $\frac{1}{4}$
- ▷  $f(i, j) = i \cdot j \cdot (i + j) \cdot (i + 2j)$
- ▷ *Unique preharmonic function* (up to multiplicative factors)
  - 📎 Biane '92

# Martin boundary theory

## Rough description

▷ *Martin boundary*  $\partial \rightsquigarrow$  set of all harmonic functions

# Martin boundary theory

## Rough description

- ▷ *Martin boundary*  $\partial \rightsquigarrow$  set of all harmonic functions
- ▷ *Minimal Martin boundary*  $\partial_m \rightsquigarrow$  integral representation of all harmonic functions:  $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$

 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92



# Martin boundary theory


## Rough description

- ▷ *Martin boundary*  $\partial \rightsquigarrow$  set of all harmonic functions
- ▷ *Minimal Martin boundary*  $\partial_m \rightsquigarrow$  integral representation of all harmonic functions:  $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$

 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

## Homogeneous random processes

- ▷ Well understood  Spitzer '64; Ney & Spitzer '66

# Martin boundary theory

## Rough description

- ▷ *Martin boundary*  $\partial \rightsquigarrow$  set of all harmonic functions
- ▷ *Minimal Martin boundary*  $\partial_m \rightsquigarrow$  integral representation of all harmonic functions:  $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$



 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

## Homogeneous random processes

- ▷ Well understood  Spitzer '64; Ney & Spitzer '66

## Non-homogeneous random processes: difficult problem

- ▷ Walks related to Lie algebras  Biane '90–'92
- ▷ Quadrant walks with **drift**  $\sum_{y \in \mathbb{N}} y \cdot p(y)$   Ignatiouk-Robert '10
- ▷ Quadrant walks with **zero drift**

Partial results by  R. '14; Bouaziz, Mustapha & Sifi '15

# Martin boundary theory

## Rough description

- ▶ *Martin boundary*  $\partial \rightsquigarrow$  set of all harmonic functions
- ▶ *Minimal Martin boundary*  $\partial_m \rightsquigarrow$  integral representation of all harmonic functions:  $f = \int_{\partial_m} \{\text{Martin kernel}\} d\mu_f$



 Martin '41;

Hunt '57; Doob '59; Choquet & Deny '60; Ney & Spitzer '66; Picardello & Woess '92

## Homogeneous random processes

- ▶ Well understood  Spitzer '64; Ney & Spitzer '66

## Non-homogeneous random processes: difficult problem

- ▶ Walks related to Lie algebras  Biane '90–'92
- ▶ Quadrant walks with **drift**  $\sum_{y \in \mathbb{N}} y \cdot p(y)$   Ignatiouk-Robert '10
- ▶ Quadrant walks with **zero drift**

Partial results by  R. '14; Bouaziz, Mustapha & Sifi '15

- ▶ General RW in cones: open problem (**conjecture**: uniqueness  $\iff$  **drift** = 0)

Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant

# Asymptotics of some numbers of walks

## Asymptotic statements




▷ *Total number of walks* starting at  $(k, l)$ :

$$q(n; k, l; \mathbb{N}^2) = \#\_{\mathbb{N}^2} \{ (k, l) \xrightarrow{n} \mathbb{N}^2 \}$$
$$\sim f_1(k, l) \cdot \rho_1^n \cdot n^{\alpha_1}$$

 Not proved yet!

# Asymptotics of some numbers of walks

## Asymptotic statements



- ▷ *Total number of walks* starting at  $(k, \ell)$ :

$$q(n; k, \ell; \mathbb{N}^2) = \#\_{\mathbb{N}^2} \{(k, \ell) \xrightarrow{n} \mathbb{N}^2\} \\ \sim f_1(k, \ell) \cdot \rho_1^n \cdot n^{\alpha_1}$$

📎 Not proved yet!


- ▷ *Excursions* starting at  $(k, \ell)$ :

$$q(n; k, \ell; i, j) = \#\_{\mathbb{N}^2} \{(k, \ell) \xrightarrow{n} (i, j)\} \\ \sim f_2(k, \ell) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$

📎 Denisov & Wachtel '15

# Asymptotics of some numbers of walks

## Asymptotic statements



- ▷ *Total number of walks* starting at  $(k, l)$ :

$$q(n; k, l; \mathbb{N}^2) = \#\_{\mathbb{N}^2} \{ (k, l) \xrightarrow{n} \mathbb{N}^2 \} \\ \sim f_1(k, l) \cdot \rho_1^n \cdot n^{\alpha_1}$$


📎 Not proved yet!

- ▷ *Excursions* starting at  $(k, l)$ :


$$q(n; k, l; i, j) = \#\_{\mathbb{N}^2} \{ (k, l) \xrightarrow{n} (i, j) \} \\ \sim f_2(k, l) \cdot f'_2(i, j) \cdot \rho_2^n \cdot n^{\alpha_2}$$

📎 Denisov & Wachtel '15

## Application: random maps with bipolar orientations



© J. Bettinelli



📎 Bousquet-Mélou, Fusy & R. '17

## Random generation

**Aim:** generate efficiently a long walk (e.g., confined to a region)




A walk of length 18000





## Random generation

**Aim:** generate efficiently a long walk (e.g., confined to a region)




A walk of length 18000

### Different methods

- ▷ *Recursive method* (step-by-step construction)
- ▷ *Bijections* (if existing)      For Kreweras see  Bernardi '07
- ▷ *Rejection algorithms*  
 Bacher & Sportiello '16; Lumbroso, Mishna & Ponty '16




## Random generation

**Aim:** generate efficiently a long walk (e.g., confined to a region)





A walk of length 18000

### Different methods

- ▷ *Recursive method* (step-by-step construction)
- ▷ *Bijections* (if existing)      For Kreweras see  Bernardi '07
- ▷ *Rejection algorithms*  
 Bacher & Sportiello '16; Lumbroso, Mishna & Ponty '16
- ▷ Preharmonic functions and *Doob transform*       Fusy '16  
(Difficulty: after Doob transform, non-uniform walks)



## Potential theoretic tools

### Counting numbers are caloric functions

- ▶ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▶ Asymptotics in three quarters of plane  Mustapha '16

## Potential theoretic tools

### Counting numbers are caloric functions



- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarters of plane  Mustapha '16

### Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone*  $f_c$ :  $\Delta[f_c] = 0$ ) containing “all” the information

## Potential theoretic tools

### Counting numbers are caloric functions

- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarters of plane  Mustapha '16



### Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone*  $f_c$ :  $\Delta[f_c] = 0$ ) containing “all” the information:

- ▷  $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$  as  $n \rightarrow \infty$

## Potential theoretic tools

### Counting numbers are caloric functions

- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarters of plane  Mustapha '16



### Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone*  $f_c$ :  $\Delta[f_c] = 0$ ) containing “all” the information:

- ▷  $q(n; k, l; \mathbb{N}^2) \approx f(k, l) \cdot \rho^n \cdot n^\alpha$  as  $n \rightarrow \infty$
- ▷  $\alpha =$  homogeneity degree of  $f_c$

## Potential theoretic tools

### Counting numbers are caloric functions

- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarters of plane  Mustapha '16



### Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone*  $f_c$ :  $\Delta[f_c] = 0$ ) containing “all” the information:

- ▷  $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$  as  $n \rightarrow \infty$
- ▷  $\alpha =$  homogeneity degree of  $f_c$
- ▷  $f \sim f_c$  asymptotically

## Potential theoretic tools

### Counting numbers are caloric functions


- ▷ Asymptotics of numbers of quadrant walks (also with inhomogeneities)  D'Arco, Lacivita & Mustapha '16
- ▷ Asymptotics in three quarters of plane  Mustapha '16

### Zero drift case: classical inequalities Varopoulos '99–'09

General principle: there is a canonical function (*the réduite of the cone*  $f_c$ :  $\Delta[f_c] = 0$ ) containing “all” the information:

- ▷  $q(n; k, \ell; \mathbb{N}^2) \approx f(k, \ell) \cdot \rho^n \cdot n^\alpha$  as  $n \rightarrow \infty$
- ▷  $\alpha =$  homogeneity degree of  $f_c$
- ▷  $f \sim f_c$  asymptotically

### Non-zero drift case: Cramér's transform & ongoing work

- ▷ Works if drift with  $\leq 0$  coordinates
- ▷ Ongoing work in the remaining cases  Garbit, Mustapha & R.



Introduction & motivations


Applications in probability theory

Applications in combinatorics

Discrete harmonic functions in the quadrant

## Functional equation & Tutte's invariants

A functional equation reminiscent of the enumeration




- ▷  $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷  $K'(x, y) = xy \left\{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \right\}$
- ▷ *Kernel functional equation:*

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$



## Functional equation & Tutte's invariants

**A functional equation** reminiscent of the enumeration



- ▷  $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷  $K'(x, y) = xy \left\{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \right\}$
- ▷ *Kernel functional equation:*

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

### Definition of Tutte's invariants


- ▷ Introduced to count  $q$ -colored triangulations & planar maps  
    📖 Tutte '73; Bernardi & Bousquet-Mélou '11
- ▷ Define  $X_0$  &  $X_1$  by  $K'(X_0, y) = K'(X_1, y) = 0$
- ▷ Tutte's invariant: function  $I \in \mathbb{Q}[[x]]$  such that  $I(X_0) = I(X_1)$

### The sections $K'(x, 0)F(x, 0)$ & $K'(0, y)F(0, y)$ are invariants

- ▷ Evaluate the functional equation at  $X_0$  &  $X_1$
- ▷ Make the difference of the two identities

# Functional equation & Tutte's invariants

**A functional equation** reminiscent of the enumeration



- ▷  $F(x, y) = \sum_{i, j \geq 1} f(i, j) x^{i-1} y^{j-1}$
- ▷  $K'(x, y) = xy \left\{ \sum_{-1 \leq k, \ell \leq 1} p(k, \ell) x^{-k} y^{-\ell} - 1 \right\}$
- ▷ *Kernel functional equation:*

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

## Definition of Tutte's invariants

- ▷ Introduced to count  $q$ -colored triangulations & planar maps  
    📖 Tutte '73; Bernardi & Bousquet-Mélou '11
- ▷ Define  $X_0$  &  $X_1$  by  $K'(X_0, y) = K'(X_1, y) = 0$
- ▷ Tutte's invariant: function  $I \in \mathbb{Q}[[x]]$  such that  $I(X_0) = I(X_1)$

## The sections $K'(x, 0)F(x, 0)$ & $K'(0, y)F(0, y)$ are invariants

- ▷ Evaluate the functional equation at  $X_0$  &  $X_1$
- ▷ Make the difference of the two identities


## Does this characterize the sections?

## Tutte's invariants & conformal mappings

**A general theorem:** uniqueness of positive harmonic functions

$K'(x, 0)F(x, 0) = w(x)$ , *characterized by*

- ▷ Conformal mapping of a certain domain
- ▷  $w(x) = w(\bar{x})$
- ▷  $w(1) = \infty$
- ▷ Same for  $K'(0, y)F(0, y)$




## Tutte's invariants & conformal mappings

**A general theorem:** description of all harmonic functions

$K'(x, 0)F(x, 0) = \text{Poly}(w(x))$ , *charac. by*

- ▷ Conformal mapping of a certain domain
- ▷  $w(x) = w(\bar{x})$
- ▷  $w(1) = \infty$
- ▷ Same for  $K'(0, y)F(0, y)$




## Tutte's invariants & conformal mappings

**A general theorem:** description of all harmonic functions

$K'(x, 0)F(x, 0) = \text{Poly}(w(x))$ , *charac. by*


- ▷ Conformal mapping of a certain domain
- ▷  $w(x) = w(\bar{x})$
- ▷  $w(1) = \infty$
- ▷ Same for  $K'(0, y)F(0, y)$



### Going back to the SRW

$K'(x, 0)F(x, 0) = \frac{x}{4(1-x)^2}$ , *characterized by*

- ▷ Conformal mapping of the unit disc
- ▷  $w(e^{i\theta}) = w(e^{-i\theta})$
- ▷  $w(1) = \infty$
- ▷ Same for  $K'(0, y)F(0, y) = \frac{y}{4(1-y)^2}$






## Tutte's invariants & conformal mappings

**A general theorem:** description of all harmonic functions

$K'(x, 0)F(x, 0) = \text{Poly}(w(x))$ , *charac. by*


- ▷ Conformal mapping of a certain domain
- ▷  $w(x) = w(\bar{x})$
- ▷  $w(1) = \infty$
- ▷ Same for  $K'(0, y)F(0, y)$



### Going back to the SRW


$K'(x, 0)F(x, 0) = \frac{x}{4(1-x)^2}$ , *characterized by*

- ▷ Conformal mapping of the unit disc
- ▷  $w(e^{i\theta}) = w(e^{-i\theta})$
- ▷  $w(1) = \infty$
- ▷ Same for  $K'(0, y)F(0, y) = \frac{y}{4(1-y)^2}$



### Question

How deep is this *connection conformal maps/harmonic functions*?



## Example: the SRW

### A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

## Example: the SRW

### A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

### Verification of the functional equation

$$K'(x, y)F(x, y) = K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0)$$

## Example: the SRW

### A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

### Verification of the functional equation

$$\begin{aligned} K'(x, y)F(x, y) &= K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \\ &= \frac{x}{4} \times \frac{1}{(1-x)^2} + \frac{y}{4} \times \frac{1}{(1-y)^2} - 0 \times 1 \end{aligned}$$

## Example: the SRW

### A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

### Verification of the functional equation

$$\begin{aligned} K'(x, y)F(x, y) &= K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \\ &= \frac{x}{4} \times \frac{1}{(1-x)^2} + \frac{y}{4} \times \frac{1}{(1-y)^2} - 0 \times 1 \end{aligned}$$

### Tutte's invariants

$$\triangleright I(X_0) = I(X_1) \xrightarrow{X_0 X_1 = 1} I(x) = I\left(\frac{1}{x}\right) \implies I \text{ function of } x + \frac{1}{x}$$

$$\triangleright \boxed{K'(x, 0)F(x, 0) = \frac{x}{4} \frac{1}{(1-x)^2} = \frac{1}{4} \frac{1}{x + \frac{1}{x} - 2}} \text{ is an invariant}$$

## Example: the SRW

### A product-form generating function

$$f(i, j) = i \cdot j \implies F(x, y) = \sum_{i, j \geq 1} i \cdot j \cdot x^{i-1} y^{j-1} = \frac{1}{(1-x)^2(1-y)^2}$$

$$\text{Kernel: } K'(x, y) = xy \left\{ \frac{x}{4} + \frac{1}{4x} + \frac{y}{4} + \frac{1}{4y} - 1 \right\} = \frac{y(x-1)^2}{4} + \frac{x(y-1)^2}{4}$$

### Verification of the functional equation

$$\begin{aligned} K'(x, y)F(x, y) &= K'(x, 0)F(x, 0) + K'(0, y)F(0, y) - K'(0, 0)F(0, 0) \\ &= \frac{x}{4} \times \frac{1}{(1-x)^2} + \frac{y}{4} \times \frac{1}{(1-y)^2} - 0 \times 1 \end{aligned}$$

### Tutte's invariants

- ▷  $I(X_0) = I(X_1) \xrightarrow{X_0 X_1 = 1} I(x) = I\left(\frac{1}{x}\right) \implies I$  function of  $x + \frac{1}{x}$
- ▷  $K'(x, 0)F(x, 0) = \frac{x}{4} \frac{1}{(1-x)^2} = \frac{1}{4} \frac{1}{x + \frac{1}{x} - 2}$  is an invariant

### Why *this* function of $x + \frac{1}{x}$ ?

- ▷ Of order 1 in  $x + \frac{1}{x} \rightsquigarrow$  *Minimality* (conformal mappings)
- ▷  $F(1, 0) = \infty \rightsquigarrow$  *Liouville's theorem*